



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

We have  $4m'^2n'^2 = 4m^2n^2 \dots\dots (5)$ .

Adding (4) and (5),  $(m'^2 + n'^2)^2 = 4m^2n^2 \left( \frac{4m^2n^2}{(m^2 - n^2)^2} + 1 \right)$

$$\text{or } m'^2 + n'^2 = \frac{2mn(m^2 + n^2)}{m^2 - n^2} \dots\dots (6).$$

$m$  and  $n$  must be selected so as to make (3) and (6) integral.

70. Proposed by CHARLES CARROLL CROSS, Libertytown, Md.

Give methods for decomposing numbers into squares, cubes, or biquadrates, and show that  $61 \times 200^3$  is the sum of ten cube numbers, and that 844933 is the sum of eleven biquadrates in thirteen different ways. [From *The Mathematical Magazine*, Vol. II, No. 10.]

Comment by the PROPOSER.

Dr. Artemas Martin has written me that he knows of no method of separating a given number into squares, cubes and other powers, except *trial*.

I take the following numbers from *The Mathematical Magazine*.

$$61 \times 200^3 = 1^3 + 2^3 + 4^3 + 6^3 + 15^3 + 46^3 + 60^3 + 270^3 + 500^3 + 700^3.$$

$$\begin{aligned} 844933 = & 27^4 + 20^4 + 16^4 + 15^4 + 12^4 + 10^4 + 8^4 + 7^4 + 3^4 + 2^4 + 1^4, \\ & = 25^4 + 24^4 + 16^4 + 15^4 + 8^4 + 6^4 + 5^4 + 4^4 + 3^4 + 2^4 + 1^4, \\ & = 25^4 + 24^4 + 15^4 + 14^4 + 12^3 + 10^4 + 7^4 + 4^4 + 3^4 + 2^4 + 1^4, \\ & = 25^4 + 22^4 + 18^4 + 15^4 + 14^4 + 11^4 + 10^4 + 6^4 + 3^4 + 2^4 + 1^4, \\ & = 25^4 + 21^4 + 18^4 + 17^4 + 14^4 + 12^4 + 10^4 + 6^4 + 5^4 + 4^4 + 1^4, \\ & = 25^4 + 21^4 + 18^4 + 16^4 + 15^4 + 12^4 + 10^4 + 9^4 + 6^4 + 3^4 + 2^4, \\ & = 25^4 + 21^4 + 18^4 + 15^4 + 14^4 + 13^4 + 12^4 + 10^4 + 8^4 + 7^4 + 2^4, \\ & = 25^4 + 20^4 + 19^4 + 18^4 + 14^4 + 10^4 + 9^4 + 7^4 + 6^4 + 4^4 + 3^4, \\ & = 25^4 + 20^4 + 19^4 + 16^4 + 14^4 + 13^4 + 11^4 + 10^4 + 9^4 + 4^4 + 2^4, \\ & = 24^4 + 23^4 + 18^4 + 15^4 + 14^4 + 12^4 + 10^4 + 9^4 + 6^4 + 5^4 + 3^4, \\ & = 24^4 + 21^4 + 20^4 + 16^4 + 15^4 + 13^4 + 10^4 + 7^4 + 6^4 + 4^4 + 1^4, \\ & = 24^4 + 21^4 + 20^4 + 16^4 + 15^4 + 12^4 + 11^4 + 8^4 + 7^4 + 5^4 + 2^4, \\ & = 24^4 + 21^4 + 19^4 + 18^4 + 15^4 + 12^4 + 10^4 + 6^4 + 5^4 + 3^4 + 2^4. \end{aligned}$$

## MISCELLANEOUS.

64. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

How many bushels of wheat will a conical bin 8 feet in diameter at base and 12 feet high hold, if part of the bin is cut off by a plane parallel to the side and passing through the center of the base?

Solution by the PROPOSER.

Let  $ABC$  be the cone,  $DF$  the plane.

Let  $BD=c$ ,  $\tan DBC = \cot FDC = n$ ,  $DC=R$ .

$\therefore x^2 + z^2 = n^2(c-y)^2$  is the equation to the cone.

$x=ny$  is the equation to the plane.

$V$ =volume of cone cut off by plane.

The limits of  $x$  are  $ny=x_2$  and  $n(c-y)=x_1$ ; of  $y$ , 0 and  $\frac{1}{2}c$ .


$$\begin{aligned}\therefore V &= 2 \int_0^{\frac{1}{2}c} \int_{x_2}^{x_1} \sqrt{[n^2(c-y)^2 - x^2]} dy dx \\ &= \int_0^{\frac{1}{2}c} \left\{ \frac{1}{2} \pi n^2 (c-y)^2 - n^2 (c-y)^2 \sin^{-1} \left( \frac{y}{c-y} \right) \right. \\ &\quad \left. - ny \sqrt{[n^2 (c-y)^2 - n^2 y^2]} \right\} dy \\ &= \frac{1}{6} \pi n^2 c^3 - \frac{2}{3} n^2 c^3 = \frac{1}{18} n^2 c^3 (3\pi - 4) \\ &= \frac{1}{18} R^2 c (3\pi - 4).\end{aligned}$$

But  $c=12$ ,  $R=4$ .

$$\therefore V = 32\pi - 1\frac{2}{3}.$$

Volume of cone  $= \frac{1}{3} \pi R^2 c = 64\pi$ .

$$\begin{aligned}\therefore \text{Required vol.} &= 64\pi - (32\pi - 1\frac{2}{3}) = 32\pi + 1\frac{2}{3} \\ &= 143.1978 \text{ cubic feet,} \\ &= 115.07 \text{ bushels.}\end{aligned}$$



Also solved by P. S. BERG and C. C. CROSS.

[NOTE.—In the figure, the point  $E$  should coincide with  $D$ . ED. F.]

65. Proposed by F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Mechanicsburg, Pa.

Show that the path of a projectile moving with a constant velocity is an inverted catenary of equal strength.

No solution has yet been received.

## PROBLEMS FOR SOLUTION.

### ARITHMETIC.

102. Proposed by ALOIS F. KOVARIK, Professor of Mathematics, Decorah Institute, Decorah, Iowa.

A's age is to B's as 2:3. 20 years from now their ages will be to each other as 4:5. What are their ages, respectively?

103. Proposed by WALTER H. DRANE, Graduate Student, Harvard University, 65 Hammond Street, Cambridge, Mass.

Find proceeds of a note discounted at a bank for 10 years at 10%. What is the meaning of the result?

\*\*\* Solutions of these problems should be sent to B. F. Finkel not later than January 10.

### ALGEBRA.

92. Proposed by ELMER SCHUYLER, High Bridge, N. J.

Given  $x^2 - yz = 1$ ;  $y^2 - xz = 2$ ;  $z^2 - xy = 3$ . Find  $x$ ,  $y$ , and  $z$ .